

NO-A186 025

STRONG CONSISTENCY OF ESTIMATION OF NUMBER OF

1/1

REGRESSION VARIABLES WHEN T (U) PITTSBURGH UNIV PA

CENTER FOR MULTIVARIATE ANALYSIS Y WU JUN 87 TR-87-15


UNCLASSIFIED

AFOSR-TR-87-1245 F49620-85-C-0008

F/G 12/3

NL





ROCOPY RESOLUTION TEST CHART

unclassified

FILE COPY

2

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD-A186 025

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR- 87-1245</b>	2. GOVT ACCESSION NO	3. RECIPIENT'S CATALOG NUMBER
TITLE (and Subtitle) Strong consistency of estimation of number of regression variables when the errors are independent and their expectations are not equal to each other.		5. TYPE OF REPORT & PERIOD COVERED <i>Final</i> <del>Technical</del> - June 1987
AUTHOR(s) Yuehua Wu		6. PERFORMING ORG. REPORT NUMBER 87-15
PERFORMING ORGANIZATION NAME AND ADDRESS Center for Multivariate Analysis University of Pittsburgh, 515 Thackeray Hall Pittsburgh, PA 15260		8. CONTRACT OR GRANT NUMBER(s) F49620-85-C-0008
CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research Department of the Air Force Bolling Air Force Base, DC 20332 nm		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 611021F 2304 A5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Same as !!		12. REPORT DATE June 1987
		13. NUMBER OF PAGES 22
		15. SECURITY CLASS. (of this report) unclassified
		16. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) DTIC ELECTE OCT 16 1987 S D		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Linear model, model selection, regression coefficient, strong consistency		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Consider the linear regression model $y_i = x_i' \beta + e_i$ , $i = 1, 2, \dots$ , where $\{x_i\}$ is a sequence of known $p$ -vectors, $\beta' = (\beta_1, \dots, \beta_p)$ is an unknown $p$ -vector, known as regression coefficients, $\{e_i\}$ is a sequence of random errors. It is of interest to test the hypothesis $H_k: \beta_{k+1} = \dots = \beta_p = 0$ , $k = 0, 1, \dots, p$ . We do not assume that the random errors are identically distributed and have zero means, since it is sometimes unrealistic. As a compensation for this relaxation, we assume the errors have a common bounded support $[a_1, a_2]$ . Under certain		

unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

conditions, we obtain the strongly consistent estimate of the number of  $k$  for which  $\beta_k \neq 0$  and  $\beta_{k+1} = \dots \beta_p = 0$ , by using the information theoretical criteria


**AFOSR-TR. 87-1245**

**STRONG CONSISTENCY OF ESTIMATION OF  
NUMBER OF REGRESSION VARIABLES WHEN  
THE ERRORS ARE INDEPENDENT AND THEIR  
EXPECTATIONS ARE NOT EQUAL TO EACH OTHER\***

*Yuehua Wu*

Center for Multivariate Analysis  
University of Pittsburgh

**Center for Multivariate Analysis  
University of Pittsburgh**



STRONG CONSISTENCY OF ESTIMATION OF  
NUMBER OF REGRESSION VARIABLES WHEN  
THE ERRORS ARE INDEPENDENT AND THEIR  
EXPECTATIONS ARE NOT EQUAL TO EACH OTHER\*

Yuehua Wu

Center for Multivariate Analysis  
University of Pittsburgh

June 1987

Technical Report No. 87-15



Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability Codes	
Dist	Avail and/or Special
A-1	

Center for Multivariate Analysis  
Fifth Floor, Thackeray Hall  
University of Pittsburgh  
Pittsburgh, PA 15260

\*Research partially supported by the Air Force Office of Scientific Research (AFSC) under contract F49620-85-C-0008. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

## Abstract

Consider the linear regression model  $y_i = x_i' \beta + e_i$ ,  $i = 1, 2, \dots$ , where  $\{x_i\}$  is a sequence of known  $p$ -vectors,  $\beta' = (\beta_1, \dots, \beta_p)$  is an unknown  $p$ -vector, known as regression coefficients,  $\{e_i\}$  is a sequence of random errors. It is of interest to test the hypothesis  $H_k: \beta_{k+1} = \dots = \beta_p = 0$ ,  $k = 0, 1, \dots, p$ . We do not assume that the random errors are identically distributed and have zero means, since it is sometimes unrealistic. As a compensation for this relaxation, we assume the errors have a common bounded support  $[a_1, a_2]$ . Under certain conditions, we obtain the strongly consistent estimate of the number  $k$  for which  $\beta_k \neq 0$  and  $\beta_{k+1} = \dots = \beta_p = 0$ , by using the information theoretical criteria.

## 1. Introduction

Consider the linear model

$$y_i = x_i' \beta + e_i, i=1, 2, \dots, n, \quad (1)$$

where  $x_i$ 's are experiment points,  $\beta = (\beta_1, \dots, \beta_n)'$  is the regression coefficient vector to be estimated, and  $e_i$ 's are random errors. In the usual linear regression model it is assumed that the random errors have vanishing expectations and common variance. In this case, the famous least square estimation (LSE) method plays an important role in making statistical inference upon the regression coefficient vector  $\beta$ . In the literature, there are a lot of papers concerning with the LSE and many important results are obtained (a part of work refers to [1],[2] and [3]). However the unbiasedness and consistency (even the weak one) of LSE strongly depend on the assumption that the expectations of errors are zero, and this assumption is not realistic sometimes. It is of interest to find a consistent estimates of the regression coefficients when the expectations of errors are not equal to each other. In [4] two methods for finding consistent estimates of the regression coefficient vector  $\beta$  are proposed.

The first method is to use the measure

$$Q_n(\beta) = \max_{1 \leq i \leq n} (y_i - x_i' \beta) - \min_{1 \leq i \leq n} (y_i - x_i' \beta)$$

The estimator  $\hat{\beta}_n$  of  $\beta$  is defined as the vector which minimizes  $Q_n(\beta)$ . The estimate  $\hat{\beta}_n$  is temporarily called MD estimate of  $\beta$  in [4] (the estimate based on the Maximum Difference between residuals)

The second method is to use the measure

$$\bar{Q}_n(\beta) = \max_{1 \leq i \leq n} |y_i - x_i' \beta|$$

Denote by  $\bar{\beta}_n$  the value of  $\beta$  which minimizes  $\bar{Q}_n(\beta)$ . Also,  $\bar{\beta}_n$  is temporarily called MA estimate of  $\beta$  (the estimate based on the Maximum Absolute values of residuals)



Under certain conditions, both  $\hat{\beta}_n$  and  $\bar{\beta}_n$  are shown to be strongly consistent in [4].

Now let us consider the hypotheses

$$H_k: \beta_{k+1} = \beta_{k+2} = \dots = \beta_p = 0 \text{ and } \beta_k \neq 0 \\ k = 0, 1, \dots, p-1.$$

It is of interest to determine the true model  $H_k$  by using the model selection criteria. Denote by  $\hat{\beta}_{kn} = (\hat{\beta}_{k1n}, \dots, \hat{\beta}_{kkn}, 0, \dots, 0)'$  the vector which minimizes  $Q_n(\beta)$  under the restriction  $\beta_{k+1} = \dots = \beta_p = 0$  and denoted by  $\bar{\beta}_{kn} = (\bar{\beta}_{k1n}, \dots, \bar{\beta}_{kkn}, 0, \dots, 0)'$  the vector which minimizes  $\bar{Q}_n(\beta)$  under the restriction  $\beta_{k+1} = \dots = \beta_p = 0$ . Write

$$\hat{Q}_k = Q_n(\hat{\beta}_{kn})$$

and

$$\bar{Q}_k = \bar{Q}_n(\bar{\beta}_{kn})$$

Choose a sequence of constants  $C_n$ , satisfying certain conditions which will be specified later, and define

$$\hat{R}_k = \hat{Q}_k + kC_n$$

and

$$\bar{R}_k = \bar{Q}_k + kC_n$$

Choose

$$\hat{k} = \text{ArgMin}\{\hat{R}_k : k \in \{0, \dots, p\}\}$$

and

$$\bar{k} = \text{ArgMin}\{\bar{R}_k : k \in \{0, \dots, p\}\}$$

where ArgMin denote the index which minimizes the quantities following the symbol ArgMin.

In this paper we shall consider the consistency of  $\hat{k}$  and  $\bar{k}$  to the true model  $k_0$ .

## 2. Consistency of $\hat{k}$

In this section, we make the following general assumptions:

Assumption 1. The errors  $e_i$ ,  $i = 1, 2, \dots$  are independent.

Assumption 2.  $P\{e_n \in [a_1, a_2]\} = 0$  and there is a positive constant  $\Delta$  such that for any  $\epsilon > 0$  and any  $n$ , we have

$$P\{e_n \in [a_1, a_1 + \epsilon]\} \geq \Delta\epsilon$$

and

$$P\{e_n \in [a_2 - \epsilon, a_2]\} \geq \Delta\epsilon.$$

Assumption 3. For any  $a > 0$ , there exists a positive constant  $C$  such that for any vector  $\alpha \neq 0$  it follows that

$$\#\{i \leq n, |l(x_i) - l(\alpha)| < a\} \geq Cn$$

for large  $n$ , hereafter  $l(\alpha) = \alpha/|\alpha|$

Assumption 4. There exists a positive constant  $m$  such that

$$|x_i| > m, \quad \text{for } i = 1, 2, \dots$$

Now let us estimate  $Q_n(\hat{\beta}_n)$ . Define

$$E_n^{(1)} = \{i \leq n, -x_i'(\hat{\beta}_n - \beta) > 0\}$$

$$E_n^{(2)} = \{i \leq n, x_i'(\hat{\beta}_n - \beta) > 0\}$$

Split  $S_p = \{x \in R^p: |x| = 1\}$  into  $d$  disjoint parts  $\Sigma_1, \dots, \Sigma_d$  such that  $\forall x, y \in \Sigma_j, x'y > 3/4$ . Let  $\gamma_j \in \Sigma_j, j = 1, \dots, d$ . Define  $E_n^j = \{i \leq n, \ell(x_i)' \gamma_j > 3/4\}, j = 1, \dots, d$ . By Assumption 3, there exists  $\delta_j > 0$  such that

$$\#(E_n^j) \geq \delta_j n, \quad j = 1, 2, \dots, d.$$

It is easy to see that  $-\ell(\hat{\beta}_n - \beta) \in \Sigma_j$  and  $i \in E_n^j$  implies that

$$-x_i' \ell(\hat{\beta}_n - \beta) > 0, \quad \text{i.e. } i \in E_n^{(1)}$$

and that  $\ell(\hat{\beta}_n - \beta) \in \Sigma_j$  and  $i \in E_n^j$  implies that

$$x_i' \ell(\hat{\beta}_n - \beta) > 0, \quad \text{i.e. } i \in E_n^{(2)}$$

Take  $r_n$  satisfying

$$r_n \rightarrow 0 \quad \text{and} \quad nr_n / \log n \rightarrow \infty,$$

we have

$$\begin{aligned}
P(Q_n(\hat{\beta}_n) \leq a_2 - a_1 - 2r_n) \\
\leq P(\max_{i \in E_n}^{(1)} e_i \leq a_2 - r_n) + P(\min_{i \in E_n}^{(2)} e_i \geq a_1 + r_n) \\
\leq \sum_{j=1}^d P(\max_{i \in E_n}^{(1)} e_i \leq a_2 - r_n, -\ell(\hat{\beta}_n - \beta) \in \Sigma_j) \\
+ \sum_{j=1}^d P(\min_{i \in E_n}^{(2)} e_i \geq a_1 + r_n, \ell(\hat{\beta}_n - \beta) \in \Sigma_j) \\
\leq \sum_{j=1}^d P(\max_{i \in E_n} e_i \leq a_2 - r_n, -\ell(\hat{\beta}_n - \beta) \in \Sigma_j) \\
+ \sum_{j=1}^d P(\min_{i \in E_n} e_i \geq a_1 + r_n, \ell(\hat{\beta}_n - \beta) \in \Sigma_j) \\
\leq \sum_{j=1}^d P(\max_{i \in E_n} e_i \leq a_2 - r_n) \\
+ \sum_{j=1}^d P(\min_{i \in E_n} e_i \geq a_1 + r_n) \\
\leq 2d(1 - \Delta r_n) \delta_1^n \leq 2de^{-\Delta r_n} \delta_1^n \leq 2d/n^2
\end{aligned}$$

for large  $n$ . By Borel-Cantelli Lemma we have

$$Q_n(\hat{\beta}_n) \geq a_2 - a_1 - 2r_n, \quad \text{a.s.}$$

when  $n$  is large enough.

Let  $k_0$  be the index of the true model and let  $\beta_0$  be the true parameter. Then obviously we have for  $p \geq k \geq k_0$

$$\begin{aligned}
Q_n(\hat{\beta}_n) &= Q_n(\hat{\beta}_{pn}) \leq Q_n(\hat{\beta}_{kn}) \\
&\leq Q_n(\hat{\beta}_{k_0n}) \leq Q_n(\beta_0) \leq a_2 - a_1
\end{aligned}$$

Thus

$$0 \leq Q_n(\hat{\beta}_{k_0n}) - Q_n(\hat{\beta}_{kn}) \leq 2r_n, \quad p \geq k \geq k_0$$

If we take  $C_n$  such that  $C_n \rightarrow 0$ ,  $C_n/r_n \rightarrow \infty$ , then for  $k > k_0$

$$\hat{R}_k - \hat{R}_{k_0} = (k - k_0)C_n + Q_n(\hat{\beta}_{kn}) - Q_n(\hat{\beta}_{k_0n}) > 0,$$

(2)

for all large  $n$ .

Next, we consider the case of  $k < k_0$ . Denote

$$\eta = |\beta_{k_0}| > 0$$

and define

$$E_n^+ = \{i \leq n, |\ell(x_i) + \ell(\hat{\beta}_{kn} - \beta_0)| < 1/2\}$$

$$E_n^- = \{i \leq n, |\ell(x_i) - \ell(\hat{\beta}_{kn} - \beta_0)| < 1/2\}$$

Split  $S_p$  into  $b$  disjoint parts  $\Pi_1, \dots, \Pi_b$  such that  $\forall x, y \in \Pi_j, |x - y| < 1/4$ . Let  $\xi_j \in \Pi_j, j = 1, \dots, b$ . Define

$$F_n^j = \{i \leq n, |\ell(x_i) - \xi_j| < 1/4\}, \quad j = 1, \dots, b.$$

By Assumption 3, there exists  $\delta_2 > 0$  such that

$$\#(F_n^j) \geq \delta_2 n, \quad j = 1, 2, \dots, b.$$

It is easy to see that

$$-\ell(\hat{\beta}_{kn} - \beta_0) \in \Pi_j \quad \text{and} \quad i \in F_n^j$$

which implies that

$$|\ell(x_i) + \ell(\hat{\beta}_{kn} - \beta_0)| < 1/2, \quad \text{i.e. } i \in E_n^+.$$

Also,

$$\ell(\hat{\beta}_{kn} - \beta_0) \in \Pi_j \quad \text{and} \quad i \in F_n^j,$$

which implies that

$$|\ell(x_i) + \ell(\hat{\beta}_{kn} - \beta_0)| < 1/2, \quad \text{i.e. } i \in E_n^-.$$

For  $i \in E_n^+$ , we have

$$x_i'(\hat{\beta}_{kn} - \beta_0) = |x_i| |\hat{\beta}_{kn} - \beta_0| \ell(x_i)' \ell(\hat{\beta}_{kn} - \beta_0)$$

$$\leq -m\eta(1 - 1/2) = -m\eta/2.$$

Similarly for  $i \in E_n^-$ , we have

$$x_i'(\hat{\beta}_{kn} - \beta_0) \geq m\eta/2.$$

Hence

$$Q_n(\hat{\beta}_{kn}) \geq \max_{i \in E_n^+} e_i - \min_{i \in E_n^-} e_i + m\eta$$

Thus

$$P(Q_n(\hat{\beta}_{kn}) \leq a_2 - a_1 + m\eta/2)$$

$$\leq P(\max_{i \in E_n^+} e_i \leq a_2 - m\eta/4) + P(\min_{i \in E_n^-} e_i \geq a_1 + m\eta/4)$$

$$\leq \sum_{j=1}^b P(\max_{i \in E_n^+} e_i \leq a_2 - m\eta/4, -\ell(\hat{\beta}_{kn} - \beta_0) \in \Pi_j)$$

$$+ \sum_{j=1}^b P(\min_{i \in E_n^-} e_i \geq a_1 + m\eta/4, \ell(\hat{\beta}_{kn} - \beta_0) \in \Pi_j)$$

$$\leq \sum_{j=1}^b P(\max_{i \in F_n^+} e_i \leq a_2 - m\eta/4, -\ell(\hat{\beta}_{kn} - \beta_0) \in \Pi_j)$$

$$+ \sum_{j=1}^b P \left( \min_{i \in F_n^j} e_i \geq a_1 + m\eta/4, \ell(\hat{\beta}_{kn} - \beta_0) \in \Pi_j \right)$$

$$\leq \sum_{j=1}^b P \left( \max_{i \in F_n^j} e_i \leq a_2 - m\eta/4 \right)$$

$$+ \leq \sum_{j=1}^b P \left( \min_{i \in F_n^j} e_i \geq a_1 + m\eta/4 \right)$$

$$\leq 2b(1 - \Delta m\eta/4) \delta_2^n \leq 2be^{-\Delta m\eta \delta_2^n/4} \leq 2b/n^2$$

for large  $n$ . By Borel-Cantelli Lemma, we have, with probability one,

$$Q_n(\hat{\beta}_n) \geq a_2 - a_1 + m\eta/2, \quad \text{for all large } n.$$

Thus for  $k < k_0$ , we have

$$\hat{R}_k - \hat{R}_{k_0} = Q_n(\hat{\beta}_{kn}) - Q_n(\hat{\beta}_{k_0n}) - (k_0 - k)C_n$$

$$\geq m\eta/2 - (k_0 - k)C_n > 0,$$

(3)

for large  $n$ , since  $C_n \rightarrow 0$ .

(2) and (3) imply that  $\hat{k}$  is strongly consistent. Summarize the above arguments, we get the following theorem.

**Theorem 1.** Choose  $C_n$  satisfying



$$(i) C_n \rightarrow 0,$$

$$(ii) nC_n / \log n \rightarrow \infty.$$

Suppose the four Assumptions given at the beginning of this section are true, then  $\hat{k} \rightarrow k$  a.s.

Proof. Use the arguments given before. We only need to note that for any sequence of  $C_n$  satisfying (i) and (ii), we can always choose  $r_n$  such that

$$(i)' r_n / C_n \rightarrow 0,$$

$$(ii)' nr_n / \log n \rightarrow \infty.$$

Q. E. D.

### 3. Consistency of $\bar{k}$

In this section, we shall make the following general assumptions:

Assumption 1: The error  $e_i$ ,  $i = 1, 2, \dots$ , are independent,

Assumption 2:  $|a_1| < a_2$ ,  $\forall n \Rightarrow P(e_n \in [a_1, a_2]) = 0$  there is a positive constant  $\tilde{\Delta}$  such that for any  $\epsilon > 0$  and for any  $n$ , we have

$$P(e_n \in [a_2 - \epsilon, a_2]) \geq \tilde{\Delta}\epsilon.$$

Assumption 3: Same as Assumption 3 in Section 2.

Assumption 4: There exists a positive constant  $\tilde{m}$  such that

$$|x_i| > \tilde{m}, \quad \text{for } i = 1, 2, \dots$$

Now let us estimate  $\bar{Q}(\beta_n)$ . Define

$$E_n = \{i \leq n, x_i'(\beta - \bar{\beta}_n) > 0\}$$

Split  $S_p$  into  $\tilde{d}$  disjoint parts  $\tilde{\Sigma}_1, \dots, \tilde{\Sigma}_{\tilde{d}}$  such that  $\forall x, y \in \tilde{\Sigma}_j, x'y > 3/4$ . Let  $\tilde{\gamma}_j \in \tilde{\Sigma}_j, j = 1, \dots, \tilde{d}$ . Define  $\tilde{E}_n^j = \{i \leq n, \ell(x_i)' \tilde{\gamma}_j > 3/4\}, j = 1, \dots, \tilde{d}$ . By Assumption 3', there exists  $\tilde{\delta}_1 > 0$  such that

$$\#(\tilde{E}_n^j) \geq \tilde{\delta}_1 n, \quad j = 1, \dots, \tilde{d}$$

It is easy to see that

$$-\ell(\bar{\beta}_n - \beta) \in \tilde{\Sigma}_j \quad \text{and} \quad i \in \tilde{E}_n^j$$

imply that  $x_i' \ell(\beta - \bar{\beta}_n) > 0$ , i. e.  $i \in E_n$ .

Take  $r_n$  satisfying

$$r_n \rightarrow 0, \quad nr_n / \log n \rightarrow \infty$$

We have

$$P(\bar{Q}_n(\bar{\beta}_n) \leq a_2 - r_n) \leq P(\max_{i \in E_n} e_i \leq a_2 - r_n)$$

$$\leq \sum_{j=1}^{\tilde{d}} P(\max_{i \in E_n} e_i \leq a_2 - r_n, \ell(\beta - \bar{\beta}_n) \in \tilde{\Sigma}_j)$$

$$\leq \sum_{j=1}^{\tilde{d}} P(\max_{i \in E_n} e_i \leq a_2 - r_n, \ell(\beta - \bar{\beta}_n) \in \tilde{\Sigma}_j)$$

$$\begin{aligned}
&\leq \sum_{j=1}^{\tilde{d}} P\left(\max_{i \in \tilde{E}_n} e_i \leq a_2 - r_n\right) \\
&\leq \tilde{d}(1 - \tilde{\Delta} r_n)^{\tilde{\delta}_1 n} \leq \tilde{d} e^{-\tilde{\Delta} r_n \tilde{\delta}_1 n} \leq \tilde{d}/n^2
\end{aligned}$$

for large  $n$ . By Borel-Cantelli Lemma we have

$$\bar{Q}_n(\bar{\beta}_n) \geq a_2 - r_n, \quad \text{a.s.}$$

when  $n$  is large enough.

Let  $k_0$  be the index of the true model and let  $\beta_0$  be the true parameter. Then obviously we have for  $p \geq k \geq k_0$

$$\begin{aligned}
\bar{Q}_n(\bar{\beta}_n) &= \bar{Q}_n(\bar{\beta}_{pn}) \leq \bar{Q}_n(\bar{\beta}_{kn}) \\
&\leq \bar{Q}_n(\bar{\beta}_{k_0 n}) \leq \bar{Q}_n(\beta_0) \leq a_2
\end{aligned}$$

Thus

$$0 \leq \bar{Q}_n(\bar{\beta}_{k_0 n}) - \bar{Q}_n(\bar{\beta}_{kn}) \leq r_n, \quad p \geq k \geq k_0.$$

If we take  $C_n$  such that

$$C_n \rightarrow 0, \quad C_n/r_n \rightarrow \infty$$

then for  $k > k_0$

$$\bar{R}_k - \bar{R}_{k_0} = (k - k_0)c_n + \bar{Q}_n(\bar{\beta}_{kn}) - \bar{Q}_n(\bar{\beta}_{k_0n}) > 0$$

(4)

for all large  $n$ .

Next, we consider the case of  $k < k_0$ . Denote

$$\tilde{\eta} = |\beta_{k_0}| > 0$$

and define

$$\bar{E}_n = \{i \leq n, \ell(x_i) \cdot \ell(\bar{\beta}_{kn} - \beta_0) \leq -1/2\}$$

Split  $S_p$  into  $\tilde{b}$  disjoint parts  $\tilde{\Pi}_1, \dots, \tilde{\Pi}_{\tilde{b}}$ , such that  $\forall x, y \in \tilde{\Pi}_j, x \cdot y \geq 1/2$ . Let  $\tilde{\xi}_j \in \tilde{\Pi}_j, j = 1, \dots, \tilde{b}$ . Define  $\bar{F}_n^j$  as  $\bar{F}_n^j = \{i \leq n, \ell(x_i) \cdot \tilde{\xi}_j \geq 275/280\}, j = 1, \dots, \tilde{b}$ . By Assumption 3, there exists  $\tilde{\delta}_2 > 0$  such that

$$\#(\bar{F}_n^j) \geq \tilde{\delta}_2 n, \quad j = 1, \dots, \tilde{b}$$

It is easy to see that  $-\ell(\bar{\beta}_{kn} - \beta_0) \in \tilde{\Pi}_j$  and  $i \in \bar{F}_n^j$  imply that

$$\ell(x_i) \cdot \ell(\bar{\beta}_{kn} - \beta_0) \leq -1/2, \quad \text{i. e. } i \in \bar{E}_n$$

For  $i \in \bar{E}_n$ , we have

$$|x_i'(\bar{\beta}_{kn} - \beta_0)| = |x_i| |\bar{\beta}_{kn} - \beta_0| |\ell(x_i)' \ell(\bar{\beta}_{kn} - \beta_0)| \geq \tilde{m}\tilde{\eta}/2$$

Hence

$$\bar{Q}_n(\bar{\beta}_{kn}) \geq \max_{i \in \bar{E}_n} e_i + \tilde{m}\tilde{\eta}/2$$

Thus

$$P(\bar{Q}_n(\bar{\beta}_{kn}) \leq a_2 + \tilde{m}\tilde{\eta}/4)$$

$$\leq P(\max_{i \in \bar{E}_n} e_i \leq a_2 - \tilde{m}\tilde{\eta}/4)$$

$$\leq \sum_{j=1}^{\tilde{b}} P(\max_{i \in \bar{E}_n} e_i \leq a_2 - \tilde{m}\tilde{\eta}/4, -\ell(\bar{\beta}_{kn} - \beta_0) \in \tilde{\Pi}_j)$$

$$\leq \sum_{j=1}^{\tilde{b}} P(\max_{i \in \bar{F}_n^j} e_i \leq a_2 - \tilde{m}\tilde{\eta}/4, -\ell(\bar{\beta}_{kn} - \beta_0) \in \tilde{\Pi}_j)$$

$$\leq \sum_{j=1}^{\tilde{b}} P(\max_{i \in \bar{F}_n^j} e_i \leq a_2 - \tilde{m}\tilde{\eta}/4)$$

$$\leq \tilde{b} (1 - \tilde{\Delta}\tilde{m}\tilde{\eta}/4) \tilde{\delta}_2^n \leq \tilde{b}/n^2$$

for large  $n$ . By Borel-Cantelli Lemma, we have with probability one, when  $n$  large enough

$$\bar{Q}_n(\bar{\beta}_{kn}) \geq a_2 + \tilde{m}\tilde{\eta}/4.$$

Thus for  $k < k_0$ , we have

$$\begin{aligned} \bar{R}_k - \bar{R}_{k_0} &= \bar{Q}_n(\bar{\beta}_{kn}) - \bar{Q}_n(\bar{\beta}_{k_0n}) - (k_0 - k)C_n \\ &\leq \tilde{m}\tilde{\eta}/4 - (k_0 - k)C_n > 0, \end{aligned} \tag{5}$$

for large  $n$ , since  $C_n \rightarrow 0$ .

(4) and (5) proves  $\bar{k}$  is consistent. Summarize the above arguments, we get the following theorem.

**Theorem 2.** Choose  $C_n$  satisfying

$$(i) \ C_n \rightarrow 0,$$

$$(ii) \ nC_n/\log n \rightarrow \infty.$$

Suppose the four assumptions given at the beginning of this section are true, then  $\bar{k} \rightarrow k$ , a. s.

**Proof.** Use the arguments given before, we only need to notice that for any sequence of  $C_n$  satisfying (i) and (ii), we can always choose  $r_n$  such that

$$(i)' \ r_n/C_n \rightarrow 0$$

$$(ii)' \ nr_n/\log n \rightarrow \infty$$

Q.E.D.

#### 4. General Case

In this section we consider the same regression model (1). But the problem we are going to solve is to determine the subset (or the model)  $J = \{1 \leq j_1 < \dots < j_k \leq p\}$  such that  $\beta_j \neq 0$  if and only if  $j \in J$ . We make the same assumptions as given in previous sections.

Of course, we can use the procedure described in section 2 and 3 to determine the model  $J$  as follows: For each permutation  $\pi$  of  $\beta = (\beta_1, \dots, \beta_p)'$ , similarly rearranging  $(x_1, \dots, x_p)'$ , we get a new model  $M_\pi$ . Under this model, using the approach given in section 2 and 3, we obtain estimates  $\hat{k} = \hat{k}_\pi = \min_{\pi} \hat{k}_\pi$  and  $\bar{k} = \bar{k}_\pi = \min_{\pi} \bar{k}_\pi$  and let  $\hat{J}_1 = \{\hat{\pi}(1), \dots, \hat{\pi}(\hat{k})\}$  and  $\bar{J}_1 = \bar{J}_1 = \{\bar{\pi}(1), \dots, \bar{\pi}(\bar{k})\}$ , we can easily prove that, by using Theorem 1 and 2,  $\hat{J}_1 \rightarrow J$ , a. s. and  $\bar{J}_1 \rightarrow J$ , a. s.

An alternative method to estimate  $J$  is given as follows: Suppose  $T$  is a subset of  $\{1, \dots, p\}$ . Consider the model  $T$ :

$$y_n = x_n(T)' \beta(T) + e_n,$$

where  $x_j(T) = (x_{ij}, j \in T)'$  and  $\beta(T) = (\beta_j, j \in T)'$ . Let

$$Q_n(T) = \min_{\beta(T)} \{ \max_{1 \leq i \leq n} (y_i - x_i(T)' \beta(T))$$

$$- \min_{1 \leq i \leq n} (y_i - x_i(T)' \beta(T)) \}$$

and

$$\bar{Q}_n(T) = \min_{\beta(T)} \max_{1 \leq i \leq n} |y_i - x_i(T)' \beta(T)|.$$

Define

$$\hat{R}_T = Q_n(T) + \#(T) C_n$$

and

$$\bar{\hat{R}}_{J_2} = \bar{Q}(T) + \#(T) C_n$$

Choose  $\hat{J}_2$  such that

$$\hat{R}_{\hat{J}_2} = \min_T \hat{R}_T$$

and choose  $\bar{J}_2$  such that

$$\bar{R}_{\bar{J}_2} = \min_T \bar{R}_T$$

We can also prove that  $\hat{J}_2 \rightarrow J$ , a. s. and  $\bar{J}_2 \rightarrow J$ , a. s. . However, there would be too much computation involved when  $p$  is relatively large. In the first case, there are totally  $p!$  permutations whileas in the second there are  $2^p$  subsets of  $\{1, \dots, p\}$ . In light of this, we propose another approach to estimate  $J$  which only involves  $p + 1$  quantities to be computed.

Now let

$$\beta(j) = (\beta_1, \dots, \beta_{j-1}, 0, \beta_{j+1}, \dots, \beta_p)'$$

and define

$$Q_n(j) = \min_{\beta(j)} \{ \max_{1 \leq i \leq n} (y_i - x_i' \beta(j)) \\ - \min_{1 \leq i \leq n} (y_i - x_i' \beta(j)) \}$$

and



$$\bar{Q}_n(j) = \min_{\beta(j)} \max_{1 \leq i \leq n} |y_i - x_i' \beta(j)|.$$

Write

$$\hat{R}(n, j) = Q_n(j) - \hat{Q}_p - c_n$$

and

$$\bar{R}(n, j) = \bar{Q}_n(j) - \bar{Q}_p - c_n.$$

We choose

$$\hat{J}_n = \{\hat{j}_1, \dots, \hat{j}_{k_n}\} = \{j: \hat{R}(n, j) > 0\}$$

and

$$\bar{J}_n = \{\bar{j}_1, \dots, \bar{j}_{k_n}\} = \{j: \bar{R}(n, j) > 0\}$$

Then we have the following theorems.

**Theorem 3.** Under the conditions of theorem 1, we have that

$$\hat{J}_n \rightarrow J, \quad \text{a. s.}$$

where model  $J = \{i_1, \dots, i_k\}$  is the true one.

**Proof** If  $j \in J$ , by (3) with the replacement that  $k_0 = p$  and  $k = p-1$ , we have that with probability one,  $\hat{R}(n, j) > 0$  for all large  $n$ , i. e.,  $j \in \hat{J}_n$ . Hence, when  $n$

large enough,  $\hat{J}_n \rightarrow J$ . Conversely, if  $j \notin J$ , using the same argument as proving theorem 1, we have

$$\begin{aligned}\hat{R}(n, j) &= Q_n(j) - \hat{Q}_p - c_n \\ &\leq 0(\log n/n) - c_n \quad \text{a. s.}\end{aligned}$$

which together with (ii) implies that

$$\hat{R}(n, j) < 0, \quad \text{for large } n,$$

i. e.  $j \notin \hat{J}_n$  when  $n$  large enough. Therefore  $\hat{J}_n \rightarrow J$  which completes the proof of Theorem 3.

**Theorem 4.** Under the conditions of theorem 2, we have that

$$\bar{J}_n \rightarrow J, \quad \text{a. s.}$$

where model  $J = \{j_1, \dots, j_k\}$  is the true one.

Proof. If  $j \in J$ , by (5) with the replacement that  $k_0 = p$  and  $k = p-1$ , we have that with probability one,  $\bar{R}(n, j) > 0$  for all large  $n$ , i. e.,  $j \in \bar{J}_n$ . Hence, when  $n$  large enough,  $\bar{J}_n \rightarrow J$ . Conversely, if  $j \notin J$ , using the same argument as proving theorem 2, we have

$$\begin{aligned}\bar{R}(n, j) &= \bar{Q}_n(j) - \bar{Q}_p - c_n \\ &\leq 0(\log n/n) - c_n \quad \text{a. s.}\end{aligned}$$

which together with (ii) implies that

$$\bar{R}(n, j) < 0, \quad \text{for large } n,$$

i. e.  $j \notin \bar{J}_n$  when  $n$  large enough. Therefore  $\bar{J}_n \subset J$  which completes the proof of Theorem 4.

## REFERENCES

1. Drygas, H. (1976). Weak and strong consistency of the least squares estimators in regression models. *Z. Wahrsch. Verw. Gebiete*, **34**, 119-127
2. Lai, T. L., Robbins, H. and Wei, C. Z. (1979). Strong consistency of least squares estimates in multiple regression II. *Journal of Multivariate Analysis*, **9**, 343-361
3. Oberhofer, W. (1982). The Consistency of nonlinear regression minimizing the  $L_1$ -norm. *Ann. Statist.* **10**, 316-319
4. Wu, Y. (1986). On strong consistent estimates of regression coefficients when the errors are not independently and identically distributed. Tech. Report No. 86-06, Center for Multivariate Analysis, Univ. of Pittsburgh

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 87-15	2. GOVT ACCESSION NO	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Strong consistency of estimation of number of regression variables when the errors are independent and their expectations are not equal to each other.		5. TYPE OF REPORT & PERIOD COVERED technical - June 1987
7. AUTHOR(s) Yuehua Wu		6. PERFORMING ORG. REPORT NUMBER 87-15
9. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Multivariate Analysis University of Pittsburgh, 515 Thackeray Hall Pittsburgh, PA 15260		8. CONTRACT OR GRANT NUMBER(s) F49620-85-C-0008
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research Department of the Air Force Bolling Air Force Base, DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE June 1987
		13. NUMBER OF PAGES 22
		15. SECURITY CLASS. (of this report) unclassified
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		18a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Linear model, model selection, regression coefficient, strong consistency		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Consider the linear regression model $y_i = x_i' \beta + e_i$ , $i = 1, 2, \dots$ , where $\{x_i\}$ is a sequence of known $p$ -vectors, $\beta' = (\beta_1, \dots, \beta_p)$ is an unknown $p$ -vector, known as regression coefficients, $\{e_i\}$ is a sequence of random errors. It is of interest to test the hypothesis $H_k: \beta_{k+1} = \dots = \beta_p = 0$ , $k = 0, 1, \dots, p$ . We do not assume that the random errors are identically distributed and have zero means, since it is sometimes unrealistic. As a compensation for this relaxation, we assume the errors have a common bounded support $[a_1, a_2]$ . Under certain		

unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

conditions, we obtain the strongly consistent estimate of the number of  $k$  for which  $\beta_k \neq 0$  and  $\beta_{k+1} = \dots \beta_p = 0$ , by using the information theoretical criteria.

unclassified

END

DATE

FILMED

DEC.

1987